HIV TESTING PATHWAY IN PRIMARY CARE


Patient requests a HIV test

Clinician suggests HIV test

HIV risk factor or HIV in differential diagnosis

Pre-test discussion:

- 1. Explain why you are recommending an HIV test e.g. "We routinely test patients with {presenting condition} for HIV because very occasionally this can be associated with HIV and we want to make sure we don't miss anything."
- 2. Ensure the patient knows the benefits of knowing HIV status
 - a. If negative: exclusion of HIV from differential & reassurance.
 - b. If positive: effective treatment, earlier treatment = better outcome
- 3. Clarify how results will be communicated (turn around ~ 2 weeks, if urgent contact HPA lab)
- 4. Inform of 3 month incubation period.
- 5. Only verbal notification that test is being done is required.
- 6. If advice needed, contact Sexual Health and HIV medicine dept, RUH (SHRUH): **01225824558.** frequently asked questions about HIV testing



HIV POSITIVE result:

Contact Sexual Health and HIV medicine dept, RUH (SHRUH) for advice: **01225824558.**

Give result in arranged way.

Appointment can be booked at SHRUH and patient may be seen within 1 week for confirmatory HIV test, in depth discussion and baseline HIV assessment.

frequently asked questions about a positive HIV

HIV NEGATIVE result:

Give result in arranged way.

If there has been risk in preceding 3 months, repeat at appropriate time.

If ongoing risk: Consider need to discuss risk reduction/ refer to SHRUH.



Royal United Hospital, Sexual Health and HIV medicine