

### What is long-term pelvic pain?

Pelvic pain is any pain you feel in the lower abdomen or pelvis. Healthcare professionals consider pelvic pain to be long-term if:

- You experience it either constantly or intermittently for at least six months
- It happens at times other than when you have your period or sexual intercourse.

Long-term pelvic pain is common. It affects around one in six women. Long-term pelvic pain is not a diagnosis in itself but a description of a symptom.

### What could long-term pelvic pain mean for me?

How we experience pain is an individual matter and may depend upon any number of factors. Long-term pain can be very difficult to live with. It may cause you emotional, social and even economic difficulties. You may experience depression, difficulties in sleeping and a disruption to your daily routine. You may fear the worst about your pain, believing that it means you have cancer or you have a serious problem that may affect you having a baby. You may have fears that people will say your pain is 'all in the mind'. The pain is not 'all in your mind'.

The reasons for long-term pelvic pain are not always easy to diagnose. It is not always possible to treat. Women may need support in managing and coping with their pain.

Even if no reasons can be found for the pain, many women find that the quality of their lives improve when they get a better understanding of what is involved.

### What causes long-term pelvic pain?

In many cases, your healthcare professionals will not be able to identify an underlying problem or give a clear diagnosis and he or she will only be able to assure you that there is no serious medical problem. Long-term pelvic pain is often caused by a combination of physical, physiological and/or social factors, rather than a single underlying condition.

These factors included:

- endometriosis - a condition where cells of the lining of the womb (the endometrium) are found elsewhere in the body, usually in the pelvis

- adenomyosis - a condition where the endometrium is in pockets within the muscle wall of the womb
- pelvic inflammatory disease (PID) – an infection of the womb, fallopian tubes and/or pelvis
- interstitial cystitis – bladder inflammation
- musculoskeletal pain – pain in your joints, muscles, ligaments and bones
- irritable bowel syndrome (IBS)
- depression, including postnatal depression
- previous or ongoing traumatic experiences such as sexual abuse in some women
- adhesions – areas of scarred tissue that may be a result of a previous infection, endometriosis or surgery – although these are common, they do not always cause pain
- trapped or damaged nerves in the pelvic area.

For some women with long-term pelvic pain none of these factors may be found.

### What happens when I first see the doctor?

At your first appointment, you should have the chance to 'tell your story', describe the pain you have experienced and discuss your anxieties. Your doctor will take your concerns seriously and listen. By working in partnership with you, he or she will aim to identify the cause(s) of your pain. Although at times you may feel you are repeating yourself, your story is important. The way you describe your symptoms is crucial in making a diagnosis. Your doctor will probably ask you a number of questions about:

- the pattern of your pain
- what makes your pain worse or better (certain sorts of movements for example)
- whether you have noticed other problems that might be linked to the pain (intercourse, bladder, bowel or psychological symptoms, for instance).

You may be asked about aspects of your everyday life including your sleep patterns, appetite and general wellbeing. You may also be asked about how you are feeling and whether you are depressed or tearful. This is because long-term pain is known to cause depression, which in turn may make your pain worse. If your healthcare professional knows how your pain is affecting you personally, this can be taking into account with your treatment.

Your doctor should explain the various factors that can lead to long-term pelvic pain. (See section 'What causes long-term pelvic pain?')

### **What types of tests might I be offered?**

Tests do not always involve getting a report from a laboratory. Your own history and the way you describe the pattern of your pain can provide much more valuable information or results. Because of this, you may be asked to keep a 'pain diary'. This involves noting down when your pain occurs, how severe it is, how long it lasts and the things that seem to affect it. A pain diary will also help you to describe your pain and to become more aware of the ways in which it affects you.

Then, depending upon your own situation, you may be offered any of the following types of tests:

- you will probably be offered an ultrasound scan.
- you may be offered screening tests for sexually transmitted infections.
- if your pain is related to psychological, bladder or bowel symptoms, your consultant may refer you to a specialist or suggest you see your GP. If you have bowel symptoms, for example, you may be referred to a gastroenterologist who may offer you tests for irritable bowel syndrome (IBS).
- if your pain occurs on a regular basis at a specific time in your menstrual cycle, then you may be offered drugs to suppress your periods for a few months. This may help your doctor in making a diagnosis.
- you may be offered a diagnostic laparoscopy. This is a procedure carried out under general anaesthetic. It involves a small cut in the abdomen to examine your reproductive organs and look for any abnormality, problems or damage. The surgeon will insert a tiny telescope (called a laparoscope) so that your reproductive organs can be seen more clearly. As with any surgical procedure there are risks and benefits and these will be explained fully to you when you are offered a test.
- if your health professional thinks that your pain is due to a particular cause, you may be offered a treatment on a 'try it and see' basis. Such treatment could help you to avoid a diagnostic laparoscopy, which carries small but significant risks.

### **What treatment may help?**

Whatever your situation, you may be offered painkillers. If these do not help to control your pain, you may be referred to a pain management team or a specialist pelvic pain clinic. Depending on the type of your pain you may also be offered other treatment.

You should be offered treatment and advice if:

- your pain is related to your menstrual cycle and you have heavy periods
- your pain varies with movement
- you have symptoms suggestive of irritable bowel syndrome
- you may have symptoms suggestive of a sexually transmitted infection or PID.

Your doctor will provide you with full information about all treatment and options.

### **Are there any risks?**

Your doctor or specialist practitioner should give you full detailed information about the risks and benefits of any investigation, surgical procedure and treatment suggested. There are no risks associated with having an ultrasound scan.

### **Are there any alternatives?**

Depending on your circumstances, you may have a range of possible options. You may be offered a combination of two or more types of treatment, such as medical treatments with tablets or injections (for example, pain relief or hormone treatment), surgery or pain management strategies. Some people find that complementary therapies can help to manage the pain.

### **What might happen if I don't have treatment?**

Your doctor may not be able to predict what might happen for you as an individual. For many women the pain gets better with time. Most women have no serious or life-threatening problems underlying the pain. Many women find that they can cope better with their pain after they have been given a thorough explanation of the nature of the pain, including previous test results and possible causes of the pain. They can also cope better when they feel reassured that there is no serious or life-threatening disease present.

### **Is there anything else I should know?**

You should visit your doctor if you experience any of the following:

- bleeding from your rectum
- a change in your bowel habits which has lasted for more than six weeks

- any new pain after you have passed the menopause
- any unusual swelling in your abdomen
- suicidal thoughts
- excessive weight loss
- irregular vaginal bleeding, such as bleeding between periods, or vaginal bleeding after the menopause or vaginal bleeding during or after sex

No treatment can be guaranteed to work all the time for everyone.

You have the right to be fully informed about your health care and have the opportunity to share in making decisions about it. Your healthcare team should respect and take account of your wishes.

If you are not comfortable with the final diagnosis, you can ask for a second opinion.

### Other organisations

IBS network  
Unit 5, 53 Mowbray Street  
Sheffield S3 8EN

Helpline: 0114 272 3253  
(Mon-Fri, 6-8pm; Sat 1am-noon)

[info@ibsnetwork.org.uk](mailto:info@ibsnetwork.org.uk)  
[www.ibsnetwork.org.uk](http://www.ibsnetwork.org.uk)  
[www.ibsnetwork.org.uk/JuniorNetwork](http://www.ibsnetwork.org.uk/JuniorNetwork)

National Endometriosis Society  
50 Westminster Palace Gardens  
Artillery Row  
London SW1P 1RR

Helpline: 0808 808 2227

[nes@endo.org.uk](mailto:nes@endo.org.uk)  
[www.endo.org.uk](http://www.endo.org.uk)

Women's Health  
52 Featherstone Street  
London EC1Y 8RT

Helpline: 0945 125 5254  
(Mon-Fri 9.30am-1.30pm)

Minicom: 0207 490 5489

[www.womenshealthlondon.org.uk](http://www.womenshealthlondon.org.uk)

### Sources and acknowledgements

This information is based on the Royal College of Obstetricians and Gynaecologists (RCOG) guideline *The Initial Management of Chronic Pelvic Pain* (published by the RCOG in April 2005.) The guidelines contain a full list of the sources of evidence used. You can find it online at [www.rcog.or.uk/index.asp?PageID=1124](http://www.rcog.or.uk/index.asp?PageID=1124).

This information has been developed by the Patient Information Subgroup of the RCOG Guidelines and Audit Committee, with input from the Consumers' Forum and the authors of the clinical guideline. It was reviewed before we published it by women attending clinics in London, Oxford and Southampton. The final version is the responsibility of the Guidelines and Audit Committee of the RCOG.

The RCOG consents to the reproduction of this document providing full acknowledgement is made. The text of this publication may accordingly be used for printing with the addition of local information or as the basis for audiotapes or for translations into other languages. Information relating to clinical recommendations must not be changed.